

Effect of Thrombin on Maturing Human Megakaryocytes

Elisabeth Martin Cramer, Jean-Marc Massé,
Jacques P. Caen, Isabel Garcia,
Janine Breton-Gorius, Najet Debili, and
William Vainchenker

From the INSERM U. 348 and Institut des Vaisseaux et du
Sang, Hôpital Lariboisière, Paris, and INSERM U. 91,
Hôpital Henri Mondor, Créteil, France

Thrombin causes platelet activation and secretion. In some nucleated cells, it is mitogenic. In this study, we have investigated how human megakaryocytes (MKs) respond to this agonist and whether the response depends on the maturation stage. MKs were cultured from bone marrow precursors in liquid culture in the presence of normal plasma. To determine whether thrombin can activate MKs, 14-day MK cultures were incubated with thrombin for 5 minutes, and cells were studied by electron microscopy, either by standard techniques or after embedding in glycolmethacrylate for immunoelectron microscopy. Ultrastructural examination of thrombin-treated MKs revealed dramatic morphological changes reminiscent of those found in platelets, including shape change and organelle centralization that involved immature as well as mature cells. MKs were also able to secrete α -granule proteins in the dilated cisternae of the demarcation membrane system, as shown by immunogold staining for thrombospondin and glycoprotein Ib. These changes were rapid (less than 5 minutes) but despite them, MKs remained viable for more than 24 hours. To determine whether thrombin has a mitogenic activity, it was added to the culture of MKs from day 3 to day 10 of culture at concentrations varying from 0.1 to 10 U/ml. Cells were subsequently studied by a double staining technique using flow cytometry to determine MK number and ploidy. No changes were observed in these two parameters, showing that thrombin is not mitogenic for MKs at the concentrations used. In conclusion, this study confirms for human MKs previous observations made about guinea pig

MKs (Fedorko et al, Lab Invest 1977, 36:32). In addition, it demonstrates that immature MKs are able to respond to thrombin and that more mature cells can secrete α -granule proteins into the demarcation membrane system, which is in continuity with the extracellular space. This phenomenon may have implications for pathological states such as myelofibrosis formation and for megakaryopoiesis autocrine regulation. (Am J Pathol 1993, 143:1498–1508)

Megakaryocytes (MKs), the bone marrow precursors of blood platelets, are mainly located along the walls of the bone marrow sinusoids during maturation.¹ They are able to cross this barrier and enter the circulation when they become mature.² MKs give rise to platelets by fragmentation of their cytoplasm into a multitude of anucleated subunits. MKs have much in common with platelets, including receptors, plasma membrane components, and storage proteins.³ The serine protease thrombin is a multifunctional key enzyme in coagulation, formed at the site of vascular injury with various biological effects: it is a potent agonist of platelet activation, eg, it activates G proteins, increases calcium and pH levels, leads to activation of phosphokinase C, phospholipase A2, and tyrosine kinases, and inhibits adenylate cyclase;⁴ it might also play an important role in vascular repair.⁵ It was reported to be a potent mitogen of diverse cell types such as fibroblasts, lymphocytes, and mesenchymal cells,^{6,7} to be chemotactic for monocytes,⁸ and to induce the expression of P-selectin on endothelial cells⁹ and the production of platelet-derived growth factor.¹⁰ Therefore, thrombin activates platelets and several other cell types causing either changes in their functional state or mitotic events.

Recently, the functional thrombin receptor has been cloned and its messenger RNA was detected in

Supported by the Association pour la Recherche sur le Cancer (ARC).

Accepted for publication July 18, 1993.

Address reprint requests to Dr. E. Martin Cramer, Laboratoire d'Hématologie, Hôpital Lariboisière, 2, rue Ambroise Paré, 75010 Paris, France.

platelets and in the megakaryocytic cell lines, Dami and HEL, suggesting that it is also expressed in normal MKs.¹¹ In this report, we investigated the effects of thrombin on human cultured MKs. Because MKs share antigenic components with platelets including glycoprotein (GP) Ib, which can bind thrombin,¹² we investigated both the activating and mitogenic effects of thrombin on human cultured MKs.

Materials and Methods

Cell Samples

Bone marrow samples were aspirated from donors for bone marrow transplantation after informed consent. They were collected into sterile syringes containing preservative-free heparin (50 U/ml) (Roche, Paris, France). Low-density cells (<1.077 g/ml) were obtained by Ficoll-metrizoate (Biochrom KG, Berlin, Germany) gradient density centrifugation and were washed three times in α medium (Eurobio, Paris, France) containing penicillin, streptomycin, and glutamine. Adherent cell depletion was performed by incubation of the cell suspension (1×10^6 per ml) in α medium with 10% fetal calf serum (Gibco, Paisley, Scotland) for at least 1 hour on plastic culture Petri dishes. Nonadherent cells were recovered and used for cell cultures in liquid medium. All these samples were obtained in accordance with the guidelines of Hospital Ethics Committee.

Megakaryocyte Liquid Culture

Low-density cells were cultured for 12 days as described previously.¹³ Briefly, Iscove's modified Dulbecco's medium (Gibco, Paisley, Scotland) was supplemented with 1% bovine serum albumin (Sigma Chemical Co., St. Louis, MO) and 10% serum derived from plasma from patients with aplastic marrows (thrombocytopenic patients after a bone marrow transplantation). Cell cultures were performed for 12 days in 25-cm² or 75-cm² tissue culture flasks (Corning Glassware Co., New York, NY) at 37°C in a 5% CO₂ fully humidified atmosphere at 1×10^6 cells per ml. Purified human thrombin (Sigma) (0.1 U/ml and 0.5 U/ml) was added once, either at day 0, 5, 7, or 9 of culture.

Electron Microscopy (EM)

MKs from liquid culture were washed twice in the presence of prostaglandin E1 (Sigma) and apyrase,

resuspended in Tyrode's buffer containing 2.9 mmol/L CaCl₂ and exposed to α -thrombin 0.1 U for 5 minutes. The reaction was stopped, and MKs were fixed in 1% glutaraldehyde in Tyrode's buffer, washed three times in phosphate buffer, and embedded either in Epon for standard EM examination, or in glycol methacrylate for immunoelectron microscopy. For immunolabeling, thin sections were incubated facing down on microwells containing polyclonal antibodies directed against GPIb¹⁴ and/or thrombospondin¹⁵ followed by goat anti-rabbit immunoglobulins coupled to 10-nm colloidal gold (Amersham les Ullys, France). The sections were observed on a Philips CM10 electron microscope.

Megakaryocyte Ploidy Measurement

A double staining technique was used to measure the ploidy of MKs. Cells in the liquid culture system (at day 10 of culture) were fixed by 2% paraformaldehyde (Serva, Heidelberg, Germany) for 20 minutes at 4°C and washed three times in phosphate-buffered saline solution. A monoclonal antibody directed against GPIIb/IIIa (Y2/51) was used to identify MKs, followed by fluorescein-conjugated goat immunoglobulin G directed against mouse immunoglobulin (Silenus, Hawthorn, Australia). Cells were washed in phosphate-buffered saline and permeabilized by 0.002% Triton X-100 (Sigma) for 20 minutes. A propidium iodide (Sigma) solution at 50 mg/ml in isotonic sodium citrate containing 100 mg/ml RNase (Boehringer, Mannheim, Germany) was used for DNA staining. Cells were analyzed by an ATC 3000 flow cytometer (Bruker, Wissembourg, France).¹³ The cell flow rate was 1,500 cells per second. The frequency of MKs in different classes was evaluated usually on 10,000 MKs. All ploidy classes over 16N were counted together because these MKs were rare in culture.

Megakaryocyte Colony Assay

The plasma clot technique was used as previously described.^{16,17} Normal serum was used at a 10% concentration. Cells were plated at 1.5×10^5 cells per ml in triplicate in the presence or the absence of 100 U/ml recombinant interleukin-3 (IL-3) (Genetics Institute, Cambridge, MA). In the assays, thrombin was added at 0.1, 0.5, 1, and 10 U/ml. At day 12, MKs were stained *in situ* by indirect immunofluorescence, using the Y2/51 monoclonal antibody.

Each dish was entirely scanned under an inverted microscope equipped for epifluorescence.

Results

Electron Microscopy

Immature MKs

Immature MKs could be distinguished from other cells by their relatively large size, high nuclear to cytoplasmic ratio, and nuclear indentation. Some of these immature MKs were forming demarcation membranes and α -granules (Figure 1a). On the very immature cells, thrombin induced membrane ruffling, nuclear condensation, and formation of nuclear blebs (Figure 1b). Immunoelectron microscopic detection of thrombospondin and GPIb performed on MKs at an intermediate maturation stage showed that some dense material composed of released α -granule proteins, such as thrombospondin (Figure 2a) was present in the forming demarcation membrane system lined by GPIb (Figure 2b).

Mature MKs

Mature MKs were recognized by their large size and cytoplasm containing numerous demarcation membranes and α -granules (Figure 3a). After thrombin was applied to the cells, dramatic morphological changes were observed: membrane ruffling with the formation of numerous long pseudopodia (Figures 1b and 4a) and the appearance of a submembranous network of demarcation membranes (Figure 3b). The nucleus often became eccentric (Figures 3b and 4b) with centralization of the other organelles, granules, and demarcation membranes, surrounded by a dense filamentous knot resembling the one observed in activated platelets (Figure 3b). The cisternae of the demarcation membrane system became widened and filled with dense material. Simultaneously, α -granules were more seldom in the cytoplasm. In our experimental conditions (no agitation), no aggregation of mature MKs was observed. Immunolabeling for thrombospondin (Figure 5a) confirmed that the dense material filling the dilated channels of the demarcation membrane system originated from α -granules, and the observation of GPIb labeling lining them (Figure 5b) confirmed that it was indeed contained within the demarcation membranes. Alpha-granules were only occasionally found in the cytoplasm of activated MKs.

Effects of Thrombin on the Proliferation of CFU-MK and MK Endomitosis (Figure 6)

In a first set of experiments, thrombin (0.1, 0.5, 1, and 10 U/ml) was added at the onset of the cultures in the colony-forming unit-MK assay. No effect was observed on the number of MK colonies when MK colonies were grown either in the absence of any added growth factor or in the presence of IL-3 (Figure 6a).

In a second set of experiments, the effect of thrombin was tested in a liquid culture technique associated with flow cytometry, which allowed to measure the number of MKs obtained in culture from the differentiation of colony-forming unit-MK (mitosis) and the ploidy of these cultured MKs (endomitosis). Thrombin was added at several concentrations (0.1 to 10 U/ml) at different days of culture. Indeed, in this culture system, polyploidization of MKs begins at day 6 of culture, and no differences in the ploidy distribution are observed after day 9. We did not find any marked differences in the number and ploidy of MKs after thrombin addition. Five tenths, 1, and 10 U/ml of thrombin gave results which were not significantly different from 0.1 U/ml (not shown). A typical experiment is shown in Figure 6b. However, these experiments were performed three times and the differences in the number of MKs observed and the ploidy distribution as calculated by the Student *t*-test were not significant.

Discussion

In this study, we have investigated the responsiveness of human MKs to thrombin stimulation. A functional thrombin receptor has been cloned and found to be present on platelets, vascular endothelial cells, and the MK-like cell lines Dami and HEL.¹¹ It was also demonstrated that MKs were cells able to be activated and respond to various stimuli such as adenosine diphosphate, thrombin, arachidonic acid, and collagen by platelet-like morphological and metabolic changes. These include thromboxane synthesis, shape change, and cytoskeletal alterations.¹⁸⁻²⁵ Thrombin has also a potent biological mitogenic effect on other cell types such as lymphocytes,⁷ endothelial cells,²⁶⁻²⁸ fibroblasts,²⁹⁻³¹ and smooth muscle cells.³² Recently, thrombin was shown to inhibit proliferation of a human megakaryoblastic cell line MEG 01, this effect involving a possible cyclic Adenosine monophosphate-dependent mechanism.³³ Thus, in this study, both the effects of thrombin on activation and mitosis of human cultured MKs were explored.




Figure 1. a: The EM appearance of this early stage of MK maturation is characterized by a moderate cell size, high nuclear to cytoplasmic ratio, rarity of α -granules (A) and prominent nucleolus (nu). The cell surface is relatively smooth and the nucleus (N) is centrally located (Go: Golgi) (6,785 \times). b: Thrombin stimulation (0.1 U/ml) of this immature MK causes nuclear bleb formation and the appearance of pseudopods (P) on the cell surface (10,250 \times).




Figure 2. Part of a middle size maturing megakaryocyte stimulated by thrombin and 0.1 U/ml for 5 minutes. **a:** Immunolabeled for thrombospondin, a marker of the forming α -granules (A): the antigen is redistributed to some intracytoplasmic secretion vacuoles (S) of this MK of intermediate maturation stage with a high nuclear to cytoplasmic ratio (dm: demarcation membranes) (36,800 \times). **b:** Immunolabeled for GPIb: after fusion with the secretion vacuoles (S) α -granules (A) are lined by immunolabeling for GPIb (arrowheads). These labeled structures correspond to the dilated channels of the demarcation membrane (dm) system (36,800 \times).




Figure 3. a: This control mature MK appears very large with low nuclear to cytoplasm ratio and has numerous α -granules (A) and prominent demarcation membrane system (dm) whose lumen is clear. The cell surface is relatively smooth (4,700 \times). b: In some thrombin-treated (0.1 U/ml) mature MKs, an eccentricity of the nucleus (N) is observed that surrounds concentric layers: a demarcation membrane zone (dm), a ring of dense filamentous meshwork (F), and a central zone containing some dilated cisternae of the demarcation membrane system filled with secretion material (S) and a few α -granules (A) (6,200 \times).




Figure 4. a: Thrombin stimulation (0.1 U/ml) of the mature cultured MK enhances the emission of long and thin pseudopods (filopodia) (f), dilatation of the demarcation membranes and concomitant secretion (S); only a few α -granules (A) remain in the cytoplasm (7,800 \times). b: Some mature MKs display a very eccentric nucleus located at one pole of the cell, almost on the process of being shed (arrowheads) from the voluminous cytoplasm. Prominent dilatation of demarcation membranes forming secretion vacuoles (S) is observed (A: α -granules) (4,800 \times).




Figure 5. Part of a large, fully mature MK stimulated by thrombin (0.1 U/ml). **a:** Immunolabeled for the α -granule protein thrombospondin: the protein is found in the large areas of secretion (S) formed by the fusion of some α -granules with the demarcation membranes (dm) (36,800 \times). **b:** Immunolabeled for GPIIb: the large secretion areas (S) are limited by the dilated channels of the demarcation membrane system because they are lined by membrane-specific labeling (arrowheads) (36,800 \times).

On the one hand, we have shown that cultured human MKs, mature as well as immature cells, strongly react to thrombin by dramatic morphological changes, including some resembling the effects of thrombin on platelets: membrane ruffling and extensive formation of pseudopods around the cell surface and/or eccentric positioning of the nucleus at one pole of the elongated cytoplasm; evidence of contractility with a thickening of the marginal zone and centralization of organelles, including granules and dilated cisternae of the demarcation membrane system and the appearance of a central, dense filamentous knot; and secretion of the α -granule content within the demarcation membrane system. The formation of nuclear blebs in the immature MKs reminded us of the nuclei of leukemic cells from megakaryoblastic leukemia (M 7) which often exhibit this peculiar and specific abnormality.³⁴ No spreading of MKs was obtained in our experimental model, unlike what Leven and Nachmias had observed in their study;²⁵ however, they used a higher dose of thrombin (1 U/ml), and full MK spreading was only observed after a long incubation time (30 minutes). Also they studied mature bone marrow MKs instead of MK cultures from progenitors. The nature of contractile response that we observed seems to differ from one cell to another, membrane ruffling (Figures 1b and 4a) and actomyosin centralization (Figure 3b). It is a possibility that these phenomena are sequential: first membrane ruffling, followed by contraction, secretion, and then by smoothing of the cell surface. Previous studies, performed on guinea pig MKs had shown that MKs could react to some platelet agonists by several structural modifications.^{18,25} The originality of our study is that it is the first to deal with human MKs and to show that not only mature but also immature MK cells can respond to thrombin. Also the use of immunoelectron microscopy has allowed us to show that thrombin induces a fusion of the α -granules with the demarcation membrane system and the subsequent secretion of granule content into the demarcation membrane channels which are in continuity with the extracellular space. It would be of interest to know whether this phenomenon is relevant in the development of myelofibrosis, as α -granules contain platelet-derived growth factor, which stimulates fibroblast growth and synthesis.³⁵⁻³⁷ It may also be involved in autocrine regulation of megakaryocytopoiesis by the release of platelet factor-4, β -thromboglobulin, and transforming growth factor- β , which are natural inhibitors of megakaryopoiesis.³⁸⁻⁴⁰ MKs have been demonstrated to be

present in large numbers in the central venous circulation,⁴¹ and evidence has also been advanced that they fragment into platelets in the lungs.⁴² It is therefore possible that the MK secretion properties described here may also be involved in pulmonary disease such as lung fibrosis.⁴³ Therefore, it might be of interest to try to develop means of controlling their possible activation.

On the other hand, our results fail to demonstrate any effect of thrombin on MK colony formation or MK ploidy, suggesting that thrombin has no mitogenic activity on either early or late stages of the MK differentiation at the concentrations used here. Although thrombin is able to activate MKs, it has no clear mitogenic effect on the MK lineage. Inactivation of thrombin by the added serum is unlikely because, even at low dose (0.1 U/ml), it could induce a strong activation of MKs as demonstrated by the ultrastructural changes. Moreover, IL-3 added in the same experimental conditions induced an increase of colony-forming unit-MK, demonstrating that the precursors were not beyond the stage of mitosis. The discrepancy between these two thrombin effects may be related to a late expression of the thrombin receptor at a differentiation stage where MK cells are incapable of mitosis or endomitosis. Alternatively, the transduction signal induced by the




Figure 6. a: Effect of thrombin (0.1 and 0.5 U/ml) and IL-3 on the number of MK colonies. b: Effect of thrombin (0.1 and 0.5 U/ml) on the cultured MK ploidy.

binding of thrombin to its receptor may cause functional response but no proliferation per se in some cell types such as MKs, monocytes, or endothelial cells.⁴⁴⁻⁴⁵ The mitogenic effect of thrombin may only be effective on G0 cells and not on cycling cells such as hematopoietic cells. Future work investigating the expression of the thrombin receptor during differentiation may be helpful to understand better these differences in the action of thrombin on MKs.

Acknowledgments

The authors wish to thank Prof. James White for improving the manuscript and Mrs. Monique Comas and Mrs. Muriel Kindelberger for secretarial assistance.

References

1. Lichtman MA, Chamberlain JK, Simon W, Santillo PA: Parasinusoidal location of megakaryocytes in marrow: a determinant of platelet release. *Am J Hematol* 1978, 4:303-312
2. Tavassoli M, Aoki M: Migration of entire megakaryocytes through the marrow-blood barrier. *Br J Haematol* 1981, 48:25-29
3. Breton-Gorius J, Vainchenker W: Expression of platelet proteins during the in vitro and in vivo differentiation of megakaryocytes and morphological aspect of their maturation. *Semin Hematol* 1986, 23:43-67
4. Shuman MA: Thrombin-cellular interactions. *Ann NY Acad Sci* 1986, 485:349-368
5. Kroll MH, Shafer AI: Biochemical mechanisms of platelet activation. *Blood* 1989, 74:1181-1195
6. Chen LB, Buchanan JM: Mitogenic activity of blood components. I. Thrombin and prothrombin. *Proc Natl Acad Sci USA* 1975, 72:131-135
7. Chen LB, Teng NNH, Buchanan JM: Mitogenicity of thrombin and surface alterations on mouse splenocytes. *Exp Cell Res* 1976, 101:41-46
8. Bar-Shavit R, Khan A, Wilner GD, Fenton JWII: Monocytes chemotaxis: stimulation by specific exosite region in thrombin. *Science* 1983, 220:728-731
9. Hattori R, Hamilton KK, Fugate RD, McEver RP, Sims PJ: Stimulated secretion of endothelial vWF is accompanied by rapid redistribution to the cell surface of the intracellular granule membrane protein GMP-140. *J Biol Chem* 1989, 264:7768-7771
10. Daniel TO, Gibbs VC, Milfay DF, Garavoy MR, Williams LT: Thrombin stimulated c-sis gene expression in microvascular endothelial cells. *J Biol Chem* 1986, 261:9579-9582
11. Vu TKH, Hung DT, Wheaton VI, Coughlin SR: Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* 1991, 64:1057-1068
12. Okamura T, Hasitz M, Jamieson GA: Platelet glycoprotein: interaction with thrombin and role as thrombin receptor on the platelet surface. *J Biol Chem* 1978, 253:3435-3443
13. Debili N, Hegyi E, Navarro S, Katz A, Mouthon MA, Breton-Gorius J, Vainchenker W: In vitro effect of hematopoietic growth factor on the proliferation, endoreplication and maturation of human megakaryocytes. *Blood* 1991, 77:2326-2338
14. Vainchenker W, Bouget J, Guichard J, Breton-Gorius J: Megakaryocyte colony formation from human bone marrow precursors. *Blood* 1979, 54:940-945
15. Cramer EM, Lu H, Caen JP, Soria C, Berndt MC, Tenza D: Differential redistribution of platelet glycoprotein Ib and IIb-IIIa after plasmin stimulation. *Blood* 1991, 77:694-699
16. Cramer EM, Debili N, Martin JF, Gladwin AM, Breton-Gorius J, Harrison P, Savidge GF, Vainchenker W: Uncoordinated appearance of fibrinogen compared with thrombospondin and von Willebrand factor in maturing human megakaryocytes. *Blood* 1989, 73:1123-1129
17. Debili N, Kieffer N, Nakazawa M, Guichard J, Titeux M, Cramer E, Breton-Gorius J, Vainchenker W: Expression of platelet glycoprotein Ib by cultured human megakaryocytes: ultrastructural localization and biosynthesis. *Blood* 1990, 76:368-376
18. Fedorko ME: The functional capacity of guinea pig megakaryocytes. I. Uptake of 3H-serotonin and morphologic response to stimuli for the platelet release reaction. *Lab Invest* 1977, 36:310-320
19. Fedorko ME: The functional capacity of guinea pig megakaryocytes. II. The uptake of particles and macro molecules and the effect of rabbit antiguinea pig platelet antiserum. *Lab Invest* 1977, 36:321-328
20. Worthington RE, Nakeff A: Thromboxane synthesis in megakaryocytes isolated by centrifugal elutriation. *Blood* 1981, 58:175-178
21. Nachmias VT: Platelet and megakaryocyte shape change: triggered alterations in the cytoskeleton. *Semin Hematol* 1983, 20:261-281
22. Miller JL: Characterization of the megakaryocyte secretory response: studies of continuously monitored release of endogenous ATP. *Blood* 1983, 61:967-972
23. Radley JM, Hartshorn MA, Green SL: The response of megakaryocytes with processes to thrombin. *Thromb Haemost* 1987, 58:732-736
24. Levine R, Eldor A, Hy AME, Gamliel H, Fuks Z, Volodavsky I: Megakaryocyte interaction with subendothelial extracellular matrix is associated with adhesion, platelet-like shape change, and thromboxane A2 production. *Blood* 1985, 66:570-577

25. Leven RM, Nachmias VT: Cultured megakaryocytes: changes in the cytoskeleton after ADP-induced spreading. *J Cell Biol* 1982, 92:313-323
26. Daniel TO, Gibbs VC, Milfay DF, Garavoy MR, Williams LT: Thrombin stimulates C-sis gene expression in microvascular endothelial cells. *J Biol Chem* 1986, 261: 9579-9582
27. Zetter BR, Antoniades HN: Stimulation of human vascular endothelial cell growth by a platelet-derived growth factor and thrombin. *J Supramol Struct* 11: 361-370
28. Gospodarowitz D, Brown KD, Birdwell CR, Zetter BR: Control of proliferation of human vascular endothelial cells. Characterization of the response of human umbilical vein endothelial cells to fibroblast growth factor, epidermal growth factor and thrombin. *J Cell Biol* 1978, 7:774-788
29. Kimura A, Katho O, Kuramoto A: Effect of platelet derived growth factor, epidermal growth factor and transforming growth factor β on the growth of human fibroblasts. *Br J Haematol* 1988, 69:9-12
30. Seuwen K, Magnaldo I, Pouyssegur J: Serotonin stimulates DNA synthesis in fibroblasts acting through 5-HT receptors coupled to a Gi-protein. *Nature* 1988, 335:254-256
31. Chambard JC, Paris S, L'Allemain G, Pouyssegur J: Two growth factor signalling pathways in fibroblasts distinguished by pertussis toxin. *Nature* 1987, 326: 800-803
32. Berk BC, Taubman MB, Cragoe EJ, Fenton JW II, Griendling KK: Thrombin signal transduction mechanisms in rat vascular smooth muscle-cells. Calcium and protein kinase C-dependent and independent pathway. *J Biol Chem* 1990, 265:17334-17340
33. Vittet D, Mathieu MN, Launay IM, Chevillard C: Thrombin inhibits proliferation of the human megakaryoblastic Meg-01 cell line: a possible involvement of a cyclic AMP-dependent mechanism. *J Cell Physiol* 1992, 150: 65-75
34. Breton-Gorius J: Phenotypes of blasts in acute erythroid blast and megakaryoblastic leukemia review. *Keio J of Med* 1987, 36:23-45
35. Breton-Gorius J, Bizet M, Reyes F, Dupuy E, Mear C, Vannier JP, Tron P: Myelofibrosis and acute megakaryoblastic leukemia in a child: topographic relationship between fibroblasts and megakaryocytes with an α -granule defect. *Leuk Res* 1982, 6:97-110
36. Castro-Malaspina H, Rabellino EM, Yen A, Nachman RL, Moore MAS: Human megakaryocyte stimulation of proliferation of bone marrow fibroblasts. *Blood* 1981, 57:781-787
37. McCarthy DM: Fibrosis of the bone marrow: content and cause. *Br J Haematol* 1985, 59:1-7
38. Gewirtz AM, Calabretta B, Rucinski B, Niewiarowski O, Xu WY: Inhibition of human megakaryocytopoiesis in vitro by platelet factor 4 (PF4) and a synthetic COOH-terminal PF4 peptide. *J Clin Invest* 1989, 83:1477-1486
39. Han ZC, Bellucci S, Tenza D, Caen JP: Negative regulation of human megakaryocytopoiesis by human platelet factor 4 and beta-thromboglobulin: comparative analysis in bone marrow culture from normal individuals and patients with essential thrombocythaemia and immune thrombocytopenic purpura. *Br J Haematol* 1990, 74:395-401
40. Mitjavila MT, Vinci G, Villeval JL, Kieffer N, Henri A, Testa U, Breton-Gorius J, Vainchenker W: Human platelet alpha-granules contain a non specific inhibitory of megakaryocyte formation: its relationship to type B-transforming growth factor (TGF-B). *J Cell Physiol* 1988, 134:93-100
41. Tingaard Pedersen N: Occurrence of megakaryocytes in various vessels and their retention in the pulmonary capillaries in man. Circulating megakaryocytes in blood from the inferior vena cava in adult rats. *Scand J Haematol* 1978, 21:369-375
42. Kaufman RM, Airo R, Pollack S, Crosby WH: Circulating megakaryocytes and platelet release in the lung. *Blood* 1965, 26:720-731
43. Martin JF, Slater DN, Trowbridge EA: Abnormal intrapulmonary platelet production. A possible cause of vascular and lung disease. *Lancet* 1983, 1:793-796
44. Low DA, Scott RW, Baker JB, Cunningham DD: Cells regulate their mitogenic response to thrombin through release of protease nexin. *Nature* 1982, 298:476-478
45. Carney DH, Scott D, Gordon EA, La Belle EF: Phosphoinositides in mitogenesis: neomycin inhibits thrombin-stimulated phosphoinositide turnover and initiation of cell proliferation. *Cell* 1985, 42:479-488